Domestic Violence

Treatment Plan For Batterers/Abusers

Psychiatric Evaluation
Comply With Prescribed Medications
Psychotherapy/Pastoral Counseling
52 Week Domestic Violence Treatment Program
No Contact With Victim Until Completing Treatment Program
Attend NA/CA/AA Meetings If Using Drugs/Alcohol
Request Help From Family & Friends



Treatment Can Be Successful

Resources

Suicide Helpline: 911 or 1-800-273-8255 Drug/Alcohol Helpline: 1-877-217-3903 Depression/Bipolar Helpline: 1-800-784-2433

Domestic Abuse Helpline: 1-800-273-8255

Domestic Violence Prevention for Dual Diagnosis & Partner Abuse

Drug/Alcohol Addiction-Mood Disorder-Domestic Violence



Developed by

Kenn I. Hicks, LCSW
Licensed Clinical Social Worker
Domestic Violence Counselor
(818) 788-8667 / mrh384@aol.com
Website: domesticviolenceprevention.org

1,800 Women Die of Domestic Violence Each Year A Woman is Battered Every 9 Seconds Every 26 Seconds a Woman is Raped

Ninety-two percent of the domestic violence assailants reported use of alcohol and/or drugs on the day of the assault

Almost every family has a member who suffers from drug and/or alcohol addiction

1 in 3 Americans could be diagnosed with a mental disorder in any given year


Dual Diagnosis

- **Dual Diagnosis:** This term is used for a mood disorder (depression/bipolar) and problems with drugs/alcohol. The person has two separate illnesses and each needs its own treatment plan.
- Partner Abuse/Domestic Violence: Along with a drug/alcohol addiction and a mood disorder, the person is a domestic violence batterer/abuser. This problem also needs its own treatment plan.

Dual Diagnosis & Partner Abuse Are All Treatable

Get Help

Drug/Alcohol Addiction



Treatment Plan

Go to a hospital or treatment center for physical and/or mental withdrawal symptoms

Psychiatric Evaluation

Attend some type of recovery group

Residential or outpatient alcohol/drug treatment program

12-Step group or another group that focuses on living without drugs and/or alcohol

Mood Disorder



Treatment Plan

Psychiatric Evaluation Comply with Prescribed Medications Psychotherapy Attend Support Group for Dual Diagnosis